



Trauma to anterior teeth in a Nigerian tertiary health facility

Dr. Osadolor OO

Department of Child Dental Health, University of Nigeria Teaching Hospital, Ituku- ozalla, Enugu State, Nigeria

Abstract

Background: Anterior teeth play a role in biting, speech and aesthetics. Trauma to anterior teeth is often neglected when not associated with pain, shocking sensation, discolouration and abscess.

Objective: To determine the distribution and pattern of presentation of trauma to anterior teeth in a tertiary health facility in South-east Nigeria.

Methodology: A retrospective review of the hospital records of patients who attended and had oral clinical evaluation in Preventive dentistry clinic of the Department of Preventive Dentistry, University of Nigeria Teaching Hospital, Enugu State, Nigeria from January 2014 to December 2015 was done.

Results: A total of 59 patients were clinically evaluated, 25(42.4%) were males, 34(57.6%) were females giving a male to female ratio of 1:1.4 The age range of the patients was 17 to 68 years with a mean age of 29.5 ± 11.8 years. Trauma to anterior teeth were more in the 20-29 age group, followed by the 30-39 age group. Enamel-dentine fractures were the most common presentation and maxillary central incisors were mostly affected. $P=0.037$, $P\text{-value}=0.031$, $P\text{-value}=0.016$.

Conclusion: The issue of delay in seeking early and proper dental treatment for traumatized anterior teeth is a common health problem. Effort needs to be applied to create better awareness on benefits of prompt and adequate treatment following trauma to anterior teeth.

Keywords: Anterior teeth, trauma, health facility

1. Introduction

Anterior teeth play a major role in biting, speech and aesthetics. Trauma to anterior teeth is often neglected when not associated with pain, shocking sensation, discolouration and abscess. Traumatic dental injuries are often due to collision with people or inanimate objects, road traffic accidents, sports, and violence [1]. These injuries if left untreated can cause, pain, psychological/ esthetics problem [2], pulpal necrosis, dental abscess and periapical pathosis, that could affect quality of life of affected individuals [3-4]. Dental injuries commonly involve anterior teeth of the upper jaw [5]. Predisposing factors of dental trauma could be related to the person's anatomic features such as increased overjet, inadequate lip coverage of the upper anterior teeth etc [6-8]. In Nigeria, the high spate of violence, increased participation in sporting activities and road traffic crashes have contributed immensely to traumatic dental injuries (TDI), thus making it an important oral health problem [9]. The issue of delay in seeking proper dental treatment is a global health problem and scientific research had shown that many factors are responsible for delay in seeking early intervention on oral health-related problems [10-19]. Some of these factors include limited access to qualified dental personnel, heavy engagements at work, ignorance, stress associated with dental visits, transportation problems, dental anxiety, ignorance, and poverty, just to mention a few [10-19]. The complications of untreated trauma to anterior teeth, may include one or a combination of the following: pulp necrosis, pulp canal obliteration, tooth discolouration, apical periodontitis, ankylosis, resorption, dento-alveolar abscess, apical granuloma or cyst [20-23]. Traumatic injuries to teeth and their supporting structures vary in severity ranging from enamel microcracks, complicated crown fracture, root

fracture to avulsions [9, 24]. The most frequent type of injury is uncomplicated crown fractures [9, 25-26]. Studies have reported traumatized teeth to be more prevalent among males [22, 24]. Single tooth involvement as the most predominant [24, 27], and maxillary incisors as being the most commonly affected [24, 28]. The most frequently traumatized teeth are the maxillary central incisors [29-31]. Road traffic accident (RTA) was reported to be the commonest cause among adults in some studies [24, 32]. Accidental damage to orodental tissues during endotracheal intubation and general anesthesia has also been reported among adults [33]. Late hospital presentation was reported as the usual practice [28, 34-35]. reported treatment given ranged from no active treatment, composite restorations to elaborate dental procedures such as root canal therapy or extraction of the affected teeth [22, 35].

Classification of diseases is the act or process of grouping diseases based on their similarities and code diagnoses, symptoms and procedures to facilitate communication among health care providers and researchers [36]. The classification of traumatic dental injuries [37] helps dental practitioners on: universal nomenclature, understanding severity of dental injuries, arriving at correct diagnosis and treatment as well as to predicting prognosis of traumatic dental injuries. Classification aids in planning the management of dental injuries and aids in describing traumatic dental injuries in epidemiological studies [36]. There are various studies and reports on trauma to anterior teeth in other parts of the country, Africa and the World. The aim of this study is to determine the distribution and pattern of presentation of trauma to anterior teeth using Ellis classification, in a tertiary health facility in South-east Nigeria. It will contribute to the existing data on trauma to

anterior teeth in Nigeria and the West African sub-region, and compare findings with published reports from Nigeria and other countries of the World.

Material and Methods

A retrospective review of the hospital records of patients who attended and had oral clinical evaluation done in Preventive dentistry clinic of the Department of Preventive Dentistry, University of Nigeria Teaching Hospital, Enugu State, Nigeria from January 2014 to December 2015 was done. University of Nigeria teaching hospital is a tertiary health facility serving many local government areas of Enugu State and neighbouring States. The demographic and clinical data, such as age, sex, diagnosis using Ellis classification and tooth/ teeth affected were retrieved from the patients' records and analyzed using SPSS version 20. Medical records with incomplete data were excluded. Ethical clearance for this study was sought from Ethical Committee of University of Nigeria Teaching Hospital, (HREC, UNTH) Enugu, and obtained before commencement. Data were analysed using a computer software programme, Statistical Package for Social Sciences (SPSS) Version 20. P values < 0.05 were accepted as being statistically significant.

Results

A total of 59 patients were clinically evaluated and Ellis classification of dental trauma was used in making the diagnosis, 25(42.4%) were males,34(57.6%) were females giving a male to female ratio of 1:1.4 The age range of the patients was 17 to 68 years with a mean age of 29.5 ± 11.8 years as shown in Table 1. Trauma to anterior teeth were seen more in the 20-29 years age group, this was followed by the 30-39 years age group as shown in Table 1. The total number of anterior teeth affected by trauma was 91 as shown in Table 2. Maxillary teeth were more affected than mandibular teeth as shown in figure 1. The maxillary central incisors were affected the most and enamel-dentine fracture were the most common presentation. Traumatized anterior teeth were more on the right side of the dental arch. P=0.037, P-value =0.031, P-value =0.016

Table 1: Socio-Demographic Characteristics OF Patients

Variable	Number	Percent
Gender		
Male	25	42.4
Female	34	57.6
Age group(yrs)		
10-19	7	11.9
20-29	31	52.5
30-39	12	20.3
40-49	4	6.8
50-59	3	5.1
60-69	2	3.4
	59	100
P-value = 0.037		

Table 2: Anterior teeth affected by trauma

Variable	Number	Percent
Central incisors	72	79.1
Lateral incisors	18	19.8
Canines	1	1.1
	91	100
P-value = 0.016		





Fig 1: Dental arch where anterior teeth were affected by trauma



P-value =0.031

Fig 2: Ellis classification of fractures seen among the affected teeth

Discussion

The mouth is the portal for life-sustaining water and food. It is also a primary mechanism for human communication [38] Anterior teeth play a major role in biting, speech and aesthetics. In this study, trauma to anterior teeth were more in the 20-29 age group, this could be due to increased physical activity in the age group, the finding was in agreement with previous studies [9,39], The correlation between age and traumatized anterior teeth was statistically significant. P-value = 0.037. Maxillary teeth were affected more than mandibular teeth, and was similar to previous reports [23, 40, 43]. Another finding in this study was that the maxillary central incisors were mostly affected by trauma, it was consistent with previous studies [2, 5, 9, 23, 43-44]. The most common type of injury was the uncomplicated crown fracture (enamel-dentine fracture) seen in (71.5%) of the traumatized teeth, similar findings were observed by many studies [2, 5, 23-26, 41-43]. The correlation between diagnosis and

Traumatized anterior teeth was statistically significant. P-value = 0.016. Females had been reported to be more concerned about their oral health and they appear to be more motivated to utilize oral health care services [45-47]. In this study, trauma to anterior teeth were seen more in females than males and was contrary to previous studies [2, 22, 24, 40, 48]. This study showed a female preponderance that could be because females have more aesthetics concerns and are more motivated to utilize oral health care services than males.

Conclusion

Trauma to anterior teeth is often neglected when not associated with pain, shocking sensation, discolouration and abscess. The issue of delay in seeking proper dental treatment is a global health problem. Effort needs to be applied to create better awareness on benefits of Immediate clinic attendance with prompt and adequate treatment following trauma to anterior teeth.

Financial support and sponsorship

None

Conflicts of interest

There are no conflicts of interest

References

- Marcenes W, Alessi ON, Traebert J. Causes and prevalence of traumatic injuries to the permanent incisors of school children aged 12 years in Jaragua do Sul, Brazil. *Int Dent J*. 2000; 50:87-92.
- Shrestha D, Upadhyay S. Pattern of Traumatic Dental Injuries and associated Risk Factors: A Hospital-based Study. *Orthodontic journal of Nepal*. 2018; 8(1):40-44.
- Glendor U. Epidemiology of traumatic dental injuries – a 12 year review of the literature. *Dental Traumatology*. 2008; 24:603–611.
- Alonge OK, Narendran S, Williamson DD. Prevalence of fractured incisal teeth among children in Harris County, Texas. *Dent Traumatol*. 2001; 17:218-21.
- Zaleckiene V, Peciuliene V, Brukiene V, Drukteinis S. Traumatic dental injuries: etiology, prevalence and possible outcomes. *Stomatologija, Baltic Dental and Maxillofacial Journal*. 2014; 16(1):7-14.
- Artun J, Behbehani F, Al-Jame B, Kerosuo H. Incisor trauma in an adolescent Arab population. Prevalence, severity and occlusal risk factors. *Am J Orthod Dentofacial Orthop*. 2005; 128:347-52.
- Traebert J, Bittencourt DD, Peres KG, Peres MA, de Lacerda JT, Marcenes W. Aetiology and rates of treatment of traumatic dental injuries among 12 years-old school children in a town in southern Brazil. *Dent Traumatol*. 2006; 22:173-8.
- Soriano EP, Caldas AF Jr, Góes PS. Risk factors related to traumatic dental injuries in Brazilian school children. *Dent Traumatol*. 2004; 20:246-50.
- Enabulele JE, Oginni AO, Sede MA, Oginni FO. Pattern of traumatized anterior teeth among adult Nigerians and complications from late presentation. *BMC Res Notes*. 2016; 9:70-79.
- Kusekwa M, Kikwilu EN. Reasons for late seeking of dental care among dental patients attending dental clinics at School of Dentistry MUHAS, Tanzania. *Tanzania Dent J*. 2011; 17(1):7–14.
- Olusile AO. Improving low awareness and inadequate access to oral health care in Nigeria: the role of dentists, the government & non-governmental agencies. *Niger Med J*. 2010; 51(3):134–6.
- Akaji EA, Oredugba FA, Jeboda SO. Utilization of dental services among secondary school students in Lagos. *Niger Dent J*. 2007; 15(2):87–91.
- Udoye CI, Oginni AO, Oginni FO. Dental anxiety among patients undergoing various dental treatments in a Nigerian teaching hospital. *J Contemp Dent Pract*. 2005; 6(2):91–8.
- Timis T, Danila I. Socio-economic status and oral health. *J Prev Med*. 2005; 13(1-2):116–21.
- Akhigbe KO, Koleoso ON. Trait anxiety, sex, age and dental treatment experience as determinants of dental anxiety among chronic dental patients in Nigeria. *Eur Sci J*. 2014; 10(12):316–28.
- Eroglu CN, Ataoglu H, Kucuk K. Factors affecting anxiety/fear of surgical procedures in dentistry. *Niger J Clin Pract*. 2017; 20(4):409-14.
- Bashiru BO, Omotunde SM. Burden of oral diseases and dental treatment needs of an urban population in Port Harcourt, Rivers State, Nigeria. *Eur J Gen Dent*. 2014; 3(2):125–8.
- Maggiras J, Locker D. Psychological factors and perceptions of pain associated with dental treatment. *Community Dent Oral Epidemiol*. 2002; 30(2):151–9.
- Patricia A, Opeyemi AB, Micah GO, Emeka CI, Adesida AA. Management of dental anxiety: A survey of Nigerian dentists. *Sahel Med J*. 2014; 17(4):159–63.
- Marcenes W, Al Beiruti N, Tayfour D, Issa S. Epidemiology of traumatic injuries to the permanent incisor of 9–12 year old school children in Damascus, Syria. *Endod Dent Traumatol*. 1999; 15:117–23.
- Saad AY. Endodontic therapy of traumatized incisors with crown and transverse intra-alveolar root fracture. *Oral Surg Oral Med Oral Pathol*. 1991; 72:468–72.
- Sandalli N, Cildir S, Guler N. Clinical investigation of traumatic injuries in Yeditepe University, Turkey during the last 3 years. *Dent Traumatol*. 2005; 21:188-94.
- Piponsuhu RA, Agbaje MO, Osisanya MO, Oyapero A. Pattern and treatment needs of traumatized anterior permanent teeth in a sub-urban area in Lagos State. *Edorium J Dent*. 2016; 3:54–62.
- Ajayi DM, Abiodun Solanke IM, Sulaiman AO, Ekhalufoh EF. A retrospective study of traumatic injuries to teeth at a Nigerian tertiary hospital. *Niger J Clin Pract*. 2012; 15:320-5.
- Caliskan MK, Turkun M. Clinical investigation of traumatic injuries of permanent incisors in Tamir, Turkey. *Endod Dent Traumatol*. 1995; 11:210–3.
- Locker D. Prevalence of traumatic dental injury in grade 8 children in six Ontario communities. *Can J Public Health*. 2005; 1:73–6.
- Zaragona AA, Catala M, Colmena ML, Valdemoro C. Dental trauma in school children six to twelve years of age. *J Dent Child*. 1998; 65:492-4.
- Zuhal K, Semra OE, Huseyin K. Traumatic injuries of the permanent incisors in children in Southern Turkey: A retrospective study. *Dent Traumatol*. 2005; 21:20-5.
- Tovo MF, dos Santos PR, Kramer PF, Feldens CA, Sari GT. Prevalence of crown fractures in 8–10 years old schoolchildren in Canoas, Brazil. *Dent Traumatol*.

- 2004; 20:251–4.
30. Rajab LD. Traumatic dental injuries in children presenting for treatment at the Department of Pediatric Dentistry, Faculty of Dentistry, University of Jordan, 1997–2000. *Dent Traumatol.* 2003; 19:6–11.
 31. Spinas E, Altana M. A new classification for crown fractures of teeth. *J Clin Pediatr Dent.* 2002; 26:225–32.
 32. Okpo EA. Fractured Permanent teeth seen in Lagos. *Niger Dent J.* 1985; 6:20-6.
 33. Skeie A, Schwartz O. Traumatic injuries of the teeth in connection with general anesthesia and the effect of use of mouth guards. *Endod Dent Traumatol.* 1999; 15:33-6.
 34. Osuji OO. Traumatized primary teeth in Nigerian children attending University Hospital: The consequences of delays in seeking treatment. *Int Dent J.* 1996; 46:165-70.
 35. Al-Jundi SH. Type of treatment, prognosis and estimation of time spent to manage dental trauma in late presentation cases at a dental teaching hospital: A longitudinal and retrospective study. *Dent Traumatol.* 2004; 20:1-5.
 36. Kikwilu EN, Minja IK, Lymo G, Mandari GJ, Kahabuka FK. Classification of traumatic dental injuries. *Dental and Maxillofacial Trauma – Challenges in Low and Middle Income Countries, 2018, 11-31.*
 37. Pagadala S, Tadikonda DC. An overview of classification of dental trauma. *IAIM, 2015; 2(9):157-164*
 38. Osadolor OO, Osadolor AJ. Oral hygiene aids in a rural community in Nigeria. *Int J Med Health Res.* 2019; 5(5):87-90.
 39. Bastone EB, Freer TJ, McNamara JR. Epidemiology of dental trauma: a review of the literature. *Aust Dent J.* 2000; 45:2–9.
 40. Zengin AZ, Celenk P, Sumer AP, Cankaya S. Evaluation of traumatic dental injuries in a group of Turkish population. *Niger J Clin Pract.* 2015; 18:86-9.
 41. Fakhruddin KS, Lawrence HP, Kenny DJ, Locker D. Etiology and environment of dental injuries in 12- to 14-year-old Ontario schoolchildren. *Dent Traumatol.* 2008; 24:305-8.
 42. Taiwo OO, Jalo HP. Dental Injuries in 12-year old Nigerian students. *Dent Traumatol.* 2011; 27:230-4.
 43. Prasad S, Tandon S, Pahuja M, Wadhawan A. Prevalence of Traumatic Dental Injuries among School Going Children In Farukhnagar, District Gurgaon. *Int J Sci Stud.* 2014; 2(2):44-49.
 44. Kirthiga M, Praveen R, Umesh W. Severity of dental trauma and its associated factors in 11-16 years old school children in Davangere city, India. *J Orofac Sci.* 2015; 7:95-9.
 45. Osadolor OO, Egbonwonu F. Pattern of demand for endodontic treatment in a Nigerian teaching hospital. *IJDMSR.* 2019; 3(6):1-4.
 46. Augusto CB, Ana HG, Cytia DE. Prevalence of endodontically treated teeth in a Brazilian Adult population. *Braz. Dent. J.* 2008; 19(4): 313-317.
 47. Omitola OG, Osagbemi T, Akadiri OA. Spectrum of diseases and pattern of referral at the Oral diagnosis clinic of a tertiary dental center. *Nig Dent J.* 2011; 19(2):66-70.
 48. Enabulele JE, Mohammed BE. Socio-Demographic Distribution of Adult Patients who Seek Oral Health

Treatment Following Trauma to Anterior Teeth J Trauma Care. 2017; 3(1):1017.